

# Polynomials and Equations

## Facts and theorems

**Theorem I (Bézout).** Let  $a$  be a root of the polynomial  $P$ . Then there exists a polynomial  $Q$  such that  $P(x)=(x-a)Q(x)$ .

**Theorem II. a)** Any polynomial of degree  $n$  has at most  $n$  roots.

**b)** If two polynomials of degree  $\leq n$  coincide in  $n+1$  points they are identical.

**Theorem III.** Any polynomial of an odd degree with real coefficients has a real root.

**Theorem IV (Viète). a)** Numbers  $a$  and  $b$  are two roots of  $x^2 + px + q = 0 \Leftrightarrow p = -(a+b), q = ab$ .

**b)** Numbers  $a, b$  and  $c$  are three roots of  $x^3+px^2+qx+r=0 \Leftrightarrow p = -(a+b+c), q = ab+ac+bc, r = -abc$ .

**Theorem V.** Given are distinct numbers  $x_1, x_2, \dots, x_n$ . Then for any numbers  $y_1, y_2, \dots, y_n$  there exist a unique polynomial  $P(x)$  of degree less than  $n$  such that  $P(x_1)=y_1, P(x_2)=y_2, \dots, P(x_n)=y_n$ .

## Problems

1. Show without algebraic calculations that the polynomials  $(x-1)(x-2)(x-3)(x-4)$  and  $(x+1)(x+2)(x+3)(x+4)$  are not identical.
2. Can the product of two nonzero polynomials be equal to the zero polynomial?
3. In the polynomial identity  $(x^2-1)(x+\dots) = (x-1)(x+3)(x+\dots)$  two numbers were replaced by dots. Restore the numbers.
4. Find the sum of all coefficients of the polynomial  $(x-2)^{100}$ .
5. Given are two polynomials  $P(x)$  and  $Q(x)$  with real coefficients. Their sets of roots are  $A$  and  $B$ , respectively. Construct a polynomial with the set of roots equal to **a)**  $A \cup B$  **b)**  $A \cap B$ .
6. How many roots has the equation  $\frac{(x-2)(x-5)}{(9-2)(9-5)} + \frac{(x-9)(x-5)}{(2-9)(2-5)} + \frac{(x-9)(x-2)}{(5-9)(5-2)} = 1$  ?
7. It is known that 10 is a root of  $x^2 + px + 2 - 5\sqrt{3} = 0$ . Find the other root of this equation.
8. Factorize the polynomials: **a)**  $x^3+x^2+x-3$  **b)**  $x^4+x^2+1$  **c)**  $x^4+1$ .
9. The product of four consecutive integers is  $7!$ . Find these integers. How many solutions has the problem?
10. Given is a polynomial  $P$ . It is known that the equation  $P(x)=x$  has no roots. Prove that the equation  $P(P(x))=x$  has no roots.

## Home work

**PE1.** It is known that the polynomials  $x^4 + ax^2 + 1$  and  $x^4 + ax^3 + 1$  have a common root. Find  $a$ .

**PE2.** Let  $P=(x+1)^{101}(x+2)^{102}(x+3)^{103} = a_{306}x^{306}+a_{305}x^{305}+\dots+a_1x+a_0$ . Find the sum  $a_0+a_2+\dots+a_{306}$ .

**PE3.** Let  $P= x^4 + x^3 - 3x^2 + x + 2$ . Prove that for any positive integer  $n>1$ , the polynomial  $P^n$  has at least one negative coefficient.

**PE4.** Given are two polynomials  $P$  and  $Q$ , both are of positive degree. It is known that the identities  $P(P(x)) \equiv Q(Q(x))$  and  $P(P(P(x))) \equiv Q(Q(Q(x)))$  hold for any  $x$ . Does that imply the identity  $P(x) \equiv Q(x)$ ?

**PE5.** For a reduced quadratic polynomial  $P$  with real coefficients, let  $M=\{x \in \mathbf{R}: |P(x)|<1\}$ . Clearly  $M$  is either empty, or is an open interval, or the union of two disjoint open intervals. Denote the sum of their lengths by  $|M|$ . Prove that  $|M|^2 \leq 8$ .